

SYPHILIS

Syphilis is a bacterial, sexually transmitted infection (STI). Syphilis is curable and progression of the disease is preventable, but if left untreated, it can cause cardiovascular and neurological diseases, blindness and can also be transmitted from a pregnant woman to her fetus.

Congenital Syphilis

is syphilis present in a pregnant mother's uterus and is acquired by her baby during birth. Women with untreated or inadequately treated early- or late-stage syphilis occurring within a year or more after infection face a 23 percent chance of an adverse pregnancy.



SYPHILIS
IF PREGNANT CAN
LEAD TO

**CONGENITAL
SYPHILIS**

CHANCE OF AN ADVERSE PREGNANCY OUTCOME IF LEFT UNTREATED.

23%

Untreated congenital syphilis can cause poor pregnancy outcomes, including miscarriages, stillbirths or death in newborns. Some infants with congenital syphilis have symptoms at birth, but many develop symptoms later.



CHANCE OF A MOTHER PASSING SYPHILIS ONTO HER UNBORN BABY IF LEFT UNTESTED.

80%

DURING THEIR FIRST
PRENATAL VISIT

1ST

EARLY THIRD
TRIMESTER

2ND

IF YOU ARE
PREGNANT
YOUR DOCTOR IS
REQUIRED TO TEST
YOU ON 3 SEPARATE
OCCASIONS

AT DELIVERY

3RD



TALK

TALK TO YOUR DOCTOR AND
ASK ABOUT SYPHILIS TESTING



TEST

GET TESTED AT YOUR 1ST PRENATAL VISIT;
DURING THE THIRD TRIMESTER NO EARLIER THAN
28 WEEKS GESTATION; AND AGAIN AT DELIVERY



TREAT

GET TREATED RIGHT AWAY IF DIAGNOSED WITH
SYPHILIS. MAKE SURE YOUR PARTNER IS TREATED, TOO



To find out more, visit MyPrenatalPromise.com



HOUSTON HEALTH
DEPARTMENT